

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01N 63/00, 27/00, C12N 7/01</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/01055</b> <b>(43) International Publication Date:</b> 18 January 1996 (18.01.96)
<b>(21) International Application Number:</b> PCT/US95/06664 <b>(22) International Filing Date:</b> 31 May 1995 (31.05.95) <b>(30) Priority Data:</b> 08/270,956      5 July 1994 (05.07.94)      US <b>(71) Applicant:</b> THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). <b>(72) Inventors:</b> HAMMOCK, Bruce, D.; 3134 Chcsapeake Bay Avenue, Davis, CA 95616 (US). McCUTCHEN, Billy, Fred; 105 Monet Circle, Willmington, DE 19808 (US). <b>(74) Agent:</b> BERLINER, Robert; Robbins, Berliner & Carson, 5th floor, 201 N. Figueroa Street, Los Angeles, CA 90012-2628 (US).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i> <i>With amended claims.</i>
<b>(54) Title:</b> INSECT CONTROL METHOD WITH GENETICALLY ENGINEERED BIOPESTICIDES  <b>(57) Abstract</b>  Insect pests can be controlled by treating the pests or their loci with a combination of recombinant virus and organic insecticide. The recombinant virus preferably is a baculovirus. Combinations of recombinant baculovirus with chemical insecticides provide a dose-response in pests, such as insects, that is greater than additive. Preferred treatments of the invention are uses of recombinant baculoviruses that express a foreign protein or toxin, with pyrethroid insecticides. Treatments against pyrethroid resistant pests are especially useful.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INSECT CONTROL METHOD WITH  
GENETICALLY ENGINEERED BIOPESTICIDES

5    Field of the Invention

          The present invention generally relates to  
uses of recombinant expression vectors expressing  
foreign proteins in controlling insects, and more  
particularly to a method utilizing insect pathogens,  
10   preferably recombinant, in conjunction with synthetic  
chemical insecticides to magnify insect kill rate.

          This invention was made with Government  
support under Grant No. 5-T32-ES07059, awarded by the  
National Institutes of Health. The Government has  
15   certain rights in this invention.

Background of the Invention

          The lepidopteran family Noctuidae includes  
some of the most destructive agricultural pests, such as  
the genera *Heliothis*, *Heliocoverpa*, *Spodoptera*, and  
20   *Trichoplusia*. For example, included in this family are  
the tobacco budworm (*Heliothis virescens*), the cotton  
bollworm (*Heliocoverpa zea*), the cotton leafworm  
(*Alabama argillacea*), the spotted cutworm (*Amathes c-*  
*nigrum*), the glassy cutworm (*Crymodes devastator*), the  
25   bronzed cutworm (*Nephelodes emmedonia*), the fall  
armyworm (*Laphygma frugiperda*), the beet armyworm  
(*Spodoptera exigua*), and the variegated cutworm  
(*Peridroma saucia*). Attempts to control these and other  
insects have often involved the use of pyrethroid

insecticides. Wildtype baculoviruses have been used with limited commercial success.

Pyrethroid insecticides now dominate the insecticide market with sales reaching into the billions annually. However, sales are beginning to stalemate due in part to the wide-scale presence of pest resistance to these compounds. In cotton alone, the presence of pyr-R *Heliothis* species has begun to result in millions of lost dollars annually. In fact, in several cases pyrethroid insecticides have completely failed to control infestations of *Heliothis* larvae in cotton, which has resulted in complete destruction of the crop. Consequently, there has been an enormous effort to control pyrethroid resistance in *Heliothis* species in cotton.

Agricultural producers sometimes attempt to restrain their use of pyrethroid insecticides until late in the growing season as a strategy against pyrethroid resistance. As a result, the producers have to turn to the less effective and more expensive organophosphate and carbamate insecticides, which have also been plagued with resistance problems. Therefore, the development of a new and effective pesticide to control pyr-R pests would be extremely valuable in any management strategy.

Resistance of agricultural pests to pesticides also leads to environmental and human health risks. These problems arise because another response by producers to pyrethroid (and other insecticide) resistance is the use of increasing amounts of pesticide, and the use of more non-selective and toxic compounds, in order to overcome pest resistance. This creates a destructive and vicious cycle.

Nevertheless, the uses of synthetic chemicals, such as exemplified by the pyrethroids, are an integral component of modern agriculture and are probably necessary to maintain our current level of agriculture

productivity, although alternative control agents, such as the earlier mentioned recombinant insect pathogens, are being explored for pest control.

Recently, the nuclear polyhedrosis virus  
5 *Autographa californica* (ACNPV), from the family Baculoviridae, has been genetically modified for an increased speed of kill by expressing insect-selective toxins. The introduction of insect-selective toxins into an insect-pathogenic virus has resulted in an  
10 approximate 30% reduction in the killing time of insect hosts, as is described by U.S. Serial No. 08/229,417, filed April 15, 1994, which is a continuation-in-part application of U.S. Serial No. 07/629,603, filed  
15 December 19, 1990, having (in part) common assignment herewith.

Among the insect specific toxins suggested for use in insect control are toxins from *Bacillus thuringiensis* from the scorpions *Buthus eupeus* and *Androctonus australis* and from the mite *Pyemotes tritici*. Further, Hammock et al., (*Nature*, 344, pp.  
20 458-461, 1990) have described use of JHE to control insects, U.S. Serial No. 07/927,851, filed August 10, 1992, and U.S. Patent 5,098,706, issued March 24, 1992, both of common assignment herewith.

25 A new tool to control resistant, as well as susceptible, insect pest populations would be very desirable.

#### Summary of the Invention

In one aspect of the present invention, a  
30 method for controlling pests is provided. Pests controlled in accordance with the invention are, for example, from the group insects, acarids, and nematodes. Such pests are treated (or their loci treated) with a synergistic combination of recombinant virus and organic  
35 insecticide. The recombinant virus preferably is a

baculovirus that expresses a foreign protein or a functional derivative thereof in pest cells infected with the recombinant baculovirus.

- 5       Treatments in accordance with the invention can be simultaneous (such as by applying a pre-mixed composition of recombinant virus and organic insecticide). Alternatively, the pests or loci may first be treated by applying virus followed by organic insecticide within about 24 hours.

10    Brief Description of the Drawing

Figure 1 illustrates "before and after" larval exposure pictures of tomato plants to illustrate a property of insect control in conjunction with the use of recombinant baculoviruses.

15    Detailed Description of the Preferred Embodiments

- 20       The present invention is the use of genetically engineered insect viruses in combination with synthetic, chemical insecticides to treat pests such as insects. Although baculoviruses will be used throughout as an illustration, this invention can be practiced with a variety of insect viruses, including DNA and RNA viruses. Using baculovirus as one example of insect viruses, we have discovered an interaction between the use of recombinant viruses in conjunction with chemical insecticides that can have an effect greater than the single effects. We will first describe suitable recombinant baculoviruses for practicing the invention.

- 30       By "baculovirus" insecticides is meant any baculovirus of the family *Baculoviridae*, such as a nuclear polyhedrosis virus (NPV). Baculoviruses are a large group of evolutionarily related viruses, which infect only arthropods; indeed, some baculoviruses only infect insects that are pests of commercially important

agricultural and forestry crops, while others are known that specifically infect other insect pests. Because baculoviruses infect only arthropods, they pose little or no risk to humans or the environment.

5           Of the suitable DNA viruses, in addition to the *Baculoviridae* are the entomopox viruses (EPV), such as *Melolontha melonotha* EPV, *Amsacta moorei* EPV, *Locusta migratoria* EPV, *Melanoplus sanguinipes* EPV, *Schistocerca gregaria* EPV, *Aedes aegypti* EPV, and  
10 *Chironomus luridus* EPV. Other suitable DNA viruses are granulosis viruses (GV). Suitable RNA viruses include togaviruses, flaviviruses, picornaviruses, cytoplasmic polyhedrosis viruses (CPV), and the like. The subfamily of double stranded DNA viruses *Eubaculovirinae* includes  
15 two genera, NPVs and GVs, which are particularly useful for biological control because they produce occlusion bodies in their life cycle. Examples of GVs include *Cydia pomonella* GV (coddling moth GV), *Pieris brassicae* GV, *Trichoplusia ni* GV, *Artogeia rapae* GV, and *Plodia*  
20 *interpunctella* GV (Indian meal moth).

          Suitable baculoviruses for practicing this invention may be occluded or non-occluded. The nuclear polyhedrosis viruses ("NPV") are one baculovirus subgroup, which are "occluded." That is, a characteristic  
25 feature of the NPV group is that many virions are embedded in a crystalline protein matrix referred to as an "occlusion body." Examples of NPVs include *Lymantria dispar* NPV (gypsy moth NPV), *Autographa californica* MNPV, *Anagrapha falcifera* NPV (celery looper NPV),  
30 *Spodoptera littoralis* NPV, *Spodoptera frugiperda* NPV, *Heliothis armigera* NPV, *Mamestra brassicae* NPV, *Choristoneura fumiferana* NPV, *Trichoplusia ni* NPV, *Heliocoverpa zea* NPV, and *Rachiplusia ou* NPV. For field use occluded viruses are preferable due to their greater  
35 stability since the viral polyhedrin coat provides protection for the enclosed infectious nucleocapsids.

Among illustrative, useful baculoviruses in practicing this invention are those *Anagrapha falcifera*, *Anticarsia gemmatalis*, *Buzura suppressoria*, *Cydia pomonella*, *Heliocoverpa zea*, *Heliothis armigera*,  
5 *Manestia brassicae*, *Plutella xylostella*, *Spodoptera exigua*, *Spodoptera littoralis*, and *Spodoptera litura*. A particularly useful "NPV" baculovirus for practicing this invention is AcNPV, which is a nuclear polyhedrosis virus from *Autographa californica*. *Autographa cali-*  
10 *fornica* is of particular interest because various major pest species within the genera *Spodoptera*, *Trichoplusia*, and *Heliothis* are susceptible to this virus.

Such baculoviruses may be genetically engineered into more potent insecticidal forms by  
15 cloning genes encoding a foreign protein such as insect toxin into the genome of the baculovirus. Illustrative toxins are from the scorpions *Buthus eupeus* and *Androctonus australis* and from the mite *Pyemotes tritici*.

20 The expressed foreign protein (which may be a glycoprotein) preferably is an insecticidal toxin, particularly an arthropod or other invertebrate toxin, such as a scorpion toxin, a wasp toxin, a snail toxin, or a spider toxin. A useful scorpion toxin is, for  
25 example, that from *Androctonus australis*. A useful snail venom is that from the snail conotoxins (cone shell snail poisons), which the animal delivers by mouth and some individual toxins of which appear to be selective for arthropods including insects. See, for  
30 example, Olivera et al., "Diversity of *Conus* Neuropeptides," *Science*, 249:257-263 (1990).

The amino acid sequence of the excitatory toxin from *Androctonus australis* (AaIT), was determined and the sequence published in Darbon 1982. The AaIT  
35 toxin exhibits toxicity to insects, while being non-toxic to isopods and mammals.



Various other scorpion toxins such as of the Buthoid scorpion can also be used, such as LqqIT2, which is a depressive insect toxin from *Leiurus quinquestriatus quinquestriatus*. The purification method used to  
5 obtain this neurotoxin was published by Zlotkin et al., *Archives of Biochem. Biophys.*, 240:877-887 (1985).

BjIT2 is another depressive insect toxin and is from *Buthotus judaicus*. The purification has been published in Lester et al., *Biochim. Biophys. Acta*,  
10 701:370-381 (1982). BjIT2 exists in two isoforms which differ in amino acid sequence at position 15. Form 1 has isoleucine in this position while form 2 has valine.

LqhIT2 is yet another depressive insect toxin from *Leiurus quinquestriatus hebraeus* which was purified  
15 using reverse phase HPLC.

An "intermediate" toxin has also been discovered which affects insect sodium channels in a manner very similar to the effect of alpha toxins on mammalian sodium channels. This neurotoxin was derived  
20 from a yellow scorpion *Leuirus quinquestriatus hebraeus*, Buthinae, Buthidae and is called herein LqhP35. The identification and purification of this toxin was described in "Toxin to Insects Derived from the Venom of the Scorpion *Leiurus quinquestriatus hebraeus*,"  
25 published by Citan et al., *Biochemistry*, 29:5941-5947 (1990), renamed "LqhQIT."

Other toxins, purified from the venom of the chactoid scorpion, *Scorpio maurus palmatus*, can also be used. For example, SmpIT2, from the chactoid scorpion,  
30 *Scorpio maurus palmatus*, is a depressive insect toxin. Its purification is described in Lazarovici et al., *J. Biol. Chem.*, 257:8397-8404 (1982).

Still other toxins purified from the venom of the chactoid scorpion, *Scorpio maurus palmatus*, are  
35 SmpCT2 and SmpCT3, and crustacean toxins, whose purification has been described in Lazarovici, Ph.D. thesis

(1980), Hebrew University, Jerusalem, "Studies on the Composition and Action of the Venom of the Scorpion *Scorpio maurus palmatus* (Scorpionidae)."

For producing recombinant baculoviruses for the purpose of controlling insects, a secretion signal sequence is preferably included. Secretion signal sequences may be derived from proteins of bacteria, yeast, fungi, or higher eukaryotes, including both animals and plants (for examples, see Watson, *Nucl. Ac. Res.*, 12:5145-5164 (1984)). More preferred are secretion signal sequences from proteins of insect origin, for example those of cecropin B from *Hyalophora cecropia* (van Hofsten et al., *PNAS*, 82:2240-2243 (1985)), and the eclosion hormone from *Manduca sexta* (Horodyski et al., *PNAS*, 86:8123-8127 (1989)). Also preferred are the secretion signal sequences naturally associated with scorpion toxins, which can be determined by the analysis of mRNA, cDNA, or genomic DNA. More preferred is the natural secretion signal sequence of AaIT (Bougis et al., *J. Biol. Chem.*, 264:19259-19265 (1989)).

Foreign proteins, or toxins, may be expressed as functional derivatives of the toxin. A "functional derivative" of the toxin is a compound which possesses a biological activity (either functional or structural) that is substantially similar to a biological activity of the toxin. The term "functional derivative" is intended to include the "fragments," "variants," "analogues," or "chemical derivatives" of a molecule. A "fragment" of a molecule such as the toxin is meant to refer to any polypeptide subset of the molecule. A "variant" of a molecule such as the toxin is meant to refer to a molecule substantially similar in structure and function to either the entire molecule, or to a fragment thereof. A molecule is said to be "substantially similar" to another molecule if both molecules have substantially similar structures or if both

molecules possess a similar biological activity. Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if the structure of one of the molecules is not found in the other, or if the sequence of amino acid residues is not identical. An "analog" of a molecule such as the toxin is meant to refer to a molecule substantially similar in function to either the entire molecule or to a fragment thereof. As used herein, a molecule is said to be a "chemical derivative" of another molecule when it contains additional chemical moieties not normally a part of the molecule.

Such moieties may improve the molecule's solubility, absorption, biological half-life, etc. Moieties capable of mediating such effects are disclosed in Remington's Pharmaceutical Sciences (1980). Procedures for coupling such moieties to a molecule are well known in the art.

Expression of the toxin will, in general, include a promoter region sufficient to direct the initiation of RNA synthesis. One baculovirus gene is that coding for polyhedrin, since the polyhedrin protein is one of the most highly expressed eucaryotic genes known. Thus, the polyhedrin promoter is preferred, although other promoter and hybrid promoter sequences may be used.

As will be further exemplified, a particular recombinant baculovirus has been constructed from *Autographa californica* by cloning a gene encoding the insect toxin from *Androctonus australis* (Hector) into the AcNPV genome, with the toxin being expressed under polyhedrin promoter control. (Such a construction is also described by U.S. Serial No. 08/229,417, already referenced and *Bio/Technology*, 9:848-852, 1991.) We designate this particular, illustrative recombinant baculovirus embodiment as "AcAaIT," but note that some

persons in the art have utilized the designation "HIT," instead of "IT," for the toxin. (See, for example, Loret et al., *Biochemistry*, 29, pp. 1492-1501 (1990), who described several of the active neurotoxins from the scorpion as "*Androctonus australis* Hector.") We have used this recombinant baculovirus (such as in comparison with the wild-type AcNPV designated as "wtAcNPV"), as will be hereinafter more fully described. However, it should be understood that the invention is not limited to the particular recombinant baculovirus ACaAIT used to illustrate aspects of this invention, although ACaAIT is particularly preferred.

Therefore, the just described recombinant baculoviruses are one part, or aspect, of practicing this invention, while the pests treatable in accordance with the present invention are from the group insects, acarids, and nematodes. For example, illustrative insect pests that can be treated are the tobacco budworm, the cotton bollworm, the cotton leafworm, the spotted cutworm, the glassy cutworm, the bronzed cutworm, the fall armyworm, the beet armyworm, and the variegated cutworm. The other part, or aspect, of practicing this invention is using such already described recombinant baculoviruses in conjunction with chemical insecticides.

The synthetic, organic insecticides with which the present method may be practiced include: Na<sup>+</sup> channel agonists (i.e. pyrethroids), Na<sup>+</sup> channel blocking agents (i.e. pyrazolines), acetylcholinesterase inhibitors (i.e. organophosphates and carbamates), nicotinic acetylcholine binding agents (e.g. imidacloprid), gabaergic binding agents (e.g. emamectin and fipronil), octapamine agonists or antagonists (i.e. formamidines), and oxphos uncouplers (e.g. pyrrole insecticides).

As will be hereinafter further exemplified in the "Experimental" section, we have used low rates of cypermethrin (a Type II pyrethroid) or allethrin (a Type I pyrethroid) in combination with wild-type ACNPV (wtACNPV) or AcAaIT (>LC<sub>99</sub>) in larvae of *H. virescens* to illustrate the invention.

The combination of the recombinant AcAaIT (even with low rates) of either pyrethroid noted above produced a dose-response greater than potentiation. Thus, use of AcAaIT in accordance with the invention in combination with allethrin or cypermethrin resulted in 54.8 and 64.6% reductions in the LT<sub>50</sub>s, respectively, of the insect larvae when compared to wtACNPV. These data illustrate a surprising property and an advantage for the invention. That is, pyrethroids and AaIT interact in a manner which magnifies the rate of kill beyond potentiation, and are synergizing (effect greater than the algebraic sum of the single effects) the rate of kill by AcAaIT. An even more surprising result is that the resistant strain of *H. virescens* is more sensitive to the recombinant baculovirus when compared with an insecticide-susceptible strain, demonstrating the potential utility of this invention.

We believe, without being bound by theory, that a simultaneous exposure of AaIT and pyrethroid results in a synergistic interaction at the sodium channel. It appears that the toxins that display synergistic activity in combination with recombinant baculovirus can be predicted based on the mechanism of action. As expected based on their activity as acetylcholinesterase inhibitors, organophosphates and carbamates were also found to be synergistic.

Thus, in practicing the invention, pests being controlled are treated (and/or their loci treated) with a combination of recombinant baculovirus and an organic insecticide predicted to be synergistic based on its

mechanism of action. The combination may conveniently be applied for treating the pests as a single admixture.

As is well known, organic insecticides may be applied by means such as spraying, atomizing, dusting, scattering or pouring and may be formulated for such applications as powders, dusts, granulates, as well as encapsulations such as in polymer substances. When practicing this invention such conventional application means may be used. Preferably, the organic insecticide and recombinant baculovirus will be admixed in desired proportions, and may typically include inert carriers such as clay, lactose, defatted soy bean powder, and the like to assist in application.

However, it is possible to apply compositions including each component separately, by utilizing the baculovirus first then followed (preferably within about forty-eight hours) by the organic insecticide. When the baculovirus is first used (followed by organic insecticide), then the baculovirus can be applied by conventional means, such as spraying. An advantage of applying the baculovirus first is due to a "falling off" property described hereinafter in Example 3.

## EXPERIMENTAL

### EXAMPLE 1

The recombinant AcAaIT that was used in performing experiments was constructed as recombinant transfer vector pAcUW2(B)·AaIT. This is orally infectious and expresses an insect-selective toxin (AaIT), which is isolated from the scorpion *Androctonus australis*, and is under the control of the p10 promoter. This construction was earlier reported in *Bio/Technology*, 9:848-852 (Sept. 1991). Briefly, the AaIT toxin gene (sometimes also herein designated

"AaHIT") in the transfer vector, pBK283, was digested with SacI and XbaI to excise the complete AaIT gene including a signal sequence of bombyxin required for secretion. The DNA fragment was inserted into the 5 pTZ-18R plasmid between the SacI and XbaI sites. The plasmid was cleaved with SacI where a synthesized SacI-BamHI-SacI linker was inserted. The resulting plasmid contained two BamHI sites; one site near the 5' end of the toxin cDNA and the other site between the 10 stop codon, and XbaI site of the original fragment. The insertion of the linker and AaIT gene was confirmed by 1% agarose gel electrophoresis by digestion with BamHI and screening for the resultant 300 base pair fragment. The protocol for GeneClean (BioL01) was followed in 15 order to isolate the toxin fragment for insertion in the transfer vector with approximately 30% recovery of the DNA fragment.

The excised AaIT gene fragment was ligated into the BglII cloning site of the digested and 20 dephosphorylated plasmid vector pAcUW2(B), and then transformed in the JM 101 strain of *E. coli*. As a result of the ligation, both the BglII and BamHI sites were eliminated, and resultant plasmids were screened for a unique SacI site. Three SacI positive clones were 25 identified from sixty colonies using 1% agarose gel electrophoresis. The direction of insertion was confirmed by double digestion with SacI (5' end of the AaIT gene) and BamHI (within the coding sequence of the polyhedrin gene of the transfer vector). Two of three 30 recombinant transfer vectors were carrying the AaIT gene in the correct orientation and had the approximate 1.6 kbp fragment inductive of the correct orientation. The construction resulted in the AaIT cDNA sequence now inserted downstream of a duplicated p10 protein promoter 35 and upstream of the polyhedrin gene 5 producing the recombinant plasmid pAcUW2(B)·AaIT.

Isolation of Recombinant AcNPV.

*Spodoptera frugiperda* cells (Sf-9) were propagated in ExCell 400 media (JR Scientific) supplemented with 2.5% fetal bovine serum. Sf-9 cells were  
5 cotransfected by calcium precipitation with the plasmid pAcUW2(B) and polyhedrin negative AcNPV DNA. Polyhedrin-infected cells were identified and collected at 5 days post infection. The recombinant virus was plaque purified by screening for the polyhedrin positive  
10 plaques. Purification of the recombinant virus was expedited by sodium dodecyl sulfate (SDS) treatment (1%) of the cells after each plaque purification in order to eliminate polyhedrin-negative, non-recombinant virus. After purification of the recombinant virus by the  
15 initial plaque assay, individual plaques were purified in 3 successive rounds, and the resultant, pure recombinants were propagated and stored at 4°C and -80°C.

Screening the Recombinant AcUW2(B)-AaIT for Biological Activity.

20 After purification of the recombinant virus by the initial plaque assay, a total of six plaques were isolated and suspended in 500 µl of ExCell media with 2.5% fetal bovine serum. Sf-9 cells in 35 mm dishes (75% monolayer) were inoculated with 100 µL (~10<sup>6</sup> plaque  
25 forming units) of the suspension, and cells were collected 7 days post infection. The individual collections were centrifuged at 1000 g for 5 minutes, and the resulting supernatant was collected and separated from the pelleted cells. The cells were  
30 resuspended in double-distilled H<sub>2</sub>O, treated with 1% SDS, then vortexed for 5 minutes and washed 3 times. Crude estimates of the plaque forming units per ml were determined from the supernatant.



EXAMPLE 2

Detailed mortality studies were conducted with 2nd instar *H. virescens* larvae, as more fully described in the Experimental Protocol of the earlier cited paper (Bio/Technology, Vol. 9). Lethal times (LTs) were derived when 2nd instar *H. virescens* larvae were treated with the recombinant and wild-type AcNPV using a common dose of 250 polyhedrin inclusion bodies (PIBs) per plug of diet.

Table 1 shows the dose response and mortality times of second instar *H. virescens* larvae for treatments with AcUW2(B)·AaIT (AcAaIT) and wild-type AcNPV.

TABLE 1

Treatment	LD (PIBs)			LT (hours)		
	10	50	90	10	50	90
AcUW2(B)·AaIT	1.56	13.3	113	59.8 <sup>a</sup>	88.0 <sup>a</sup>	129 <sup>a</sup>
Wild-Type AcNPV	2.72	21.9	175	91.4	125	172

<sup>a</sup> Significantly different from other treatments - POLO probit analysis program (C.I. 0.95)

PIBs = polyhedrin inclusion bodies  
LT = lethal time; LD = lethal dose

These data demonstrate the ability of the AcUW2(B)·AaIT to kill *H. virescens* larvae significantly more quickly than the wild-type AcNPV.

EXAMPLE 3

Larvae infected with the recombinant AcAaIT typically started showing symptoms of paralysis and stopped feeding 10 to 15 hours prior to death. As a result, these larvae can functionally be considered dead, and treatment with AcAaIT represents an approxi-

mately 40% reduction in the time required to kill host larvae when compared to wild-type AcNPV.

We designed experiments to assess the differences in feeding damage that would be incurred by tomato plants infested with larvae of *H. virescens* that had been infected with uninfected controls wild-type AcNPV or AcAaIT. Individual plants were infested with two 4th instar larvae that were uninfected (control, I) or had previously been infected with wild-type AcNPV (II) or /AcAaIT (III) 48 hours earlier. These larvae were allowed to feed continuously on the plants until death.

The AcAaIT treatment resulted in significantly less damaged plants. However, what is not evident is the fact that the larvae infected with AcAaIT were unable to inflict more damage on the tomato plants because they were falling from the plant. In fact, we observed five of eight 4th and 5th instar larvae infected with AcAaIT had fallen from the plant at 120 hours post infection. These individuals were apparently unable to crawl back up the plant due to the onset of paralysis and is sometimes referred to as the "falling off" property. This is illustrated by Fig. 1.

In Fig. 1, panels A illustrate the tomato plants before exposure to any larvae. Panel IA thus shows the control "before" larvae while panel IB shows the control after larvae without any virus had been freely feeding on the plant until the experiment was stopped. As demonstrated by the illustrations (reproduced from photographs), the control tomato plants were badly ravaged by the larvae. Turning to panel IIA, we again see tomato plants before introduction of larvae. In panel IIB larvae infected with wild-type AcNPV have been feeding on the plants for about 190-200 hours at which point the larvae had died and the experiment (for all panels IB, IIB, and IIIB) was

stopped. While panel IIB showed considerable improvement over the panel IB control, the plant nevertheless had been subjected to up to about 200 hours of feeding activity before the larvae died. However, turning to panels IIIA (the beginning of the experiment) and IIIB (the end), we see that the panel IIIB plants have considerably more foliage when the experiment was stopped. These plants have more foliage because the inventive baculovirus AcAaIT infected larvae had their behavior disrupted before death so that five of eight larvae fell off the plant after 120 hours of feeding and the remaining three fell shortly thereafter.

Table 2 confirms that the insects become irritated and were not feeding on the food substrate or were wandering off the substrate a significant amount of time.

TABLE 2

Percentage of Time During Course of Infection  
Tobacco Budworm Larvae Were Off the Food Substrate

<u>Food Substrate</u>	<u>AcAaIT</u>	<u>Wild Type Virus</u>
Cotton	80.40%	13.30%
Romaine	64.20%	15.40%
Iceberg	85.90%	13.20%
Diet	59.90%	11.00%

To obtain the data summarized by Table 2, the larvae were droplet fed virus at 2,000 PIBs/ $\mu$ l as neonates and checked every 6 to 8 hours. Larvae were scored as to whether they were off or on the food substrate as summarized in Table 2 above. At 64 hours post-infection, over 57% of all tobacco budworm larvae were not on the substrate, versus 32% of wild-type infected insects (which were not yet expressing symptoms). The purpose of this experiment was to confirm that larvae infected with AcAaIT before they die

become paralyzed (or have their feeding behavior disrupted), while this does not happen with wild-type infected insects.

It is plausible that low levels of the neurotoxin, AaIT, early in the infection process may irritate the larval host, thus resulting in the "falling off" or eating disruption phenomena. Even though the larva is still potentially capable of feeding and causing damage, it no longer has access to the plant. With the reduced killing time and early immobilization of the host larva by paralysis, AcaAaIT appears effectively to curtail food consumption which may translate into reduced feeding damage.

#### EXAMPLE 4

The slow time to death and resulting crop damage is often cited as a major limitation to the commercial success of viral insecticides. Twenty-four to forty-eight hour mortality is a common target for classical insecticides. Thus, in this study we have concentrated on studying speed of kill expressed as lethal times ("LTs"). Analogous approaches can be used to determine lethal doses, which are likely to be of major economic importance.

As is summarized by Tables 3A and 3B, we investigated the interaction of low rates ( $LC_{10}$ - $LC_{20}$  at 24 h; Table 4) of six insecticides in combination with wild-type AcNPV or AcaAaIT ( $>LC_{99}$ ) in neonate larvae of *H. virescens*. For example, two of the compounds studied were pyrethroids, allethrin (Type I pyrethroid). In our study we found a subadditive response when low rates of allethrin or cypermethrin were combined with wild-type AcNPV. Thus, a combination of allethrin and wild-type AcNPV reduced the  $LT_{50}$  to 71.9 h (53.1 h, theoretical additive effect), which is a 17.9% reduction when

compared to wild-type AcNPV alone. Likewise, cypermethrin in combination with wild-type AcNPV produced a  $LT_{50}$  of 65.8 h (52.1 h, theoretical), which represents a 24.2% reduction when compared to wild-type AcNPV. These results suggest that the pyrethroids and the wild-type AcNPV act independently of one another, but result in a subadditive response since the combination is positive although it is significantly less than a theoretical interaction (additive effect).

5

**TABLE 3A****Insecticides Combined with Wild-Type  
and Recombinant Baculoviruses**

	<u>Compound</u>	<u>Common Name</u>	<u>Type</u>	<u>Mode of Action</u>	<u>Source</u>
5	Allethrin	Cinerine, Alleviate, Bioallethrin	Pyrethroid Type I	Sodium Channel Agonist	Roussel Uclaf Div. Agrovetenaire, Paris, France
10	Cypermethrin	Ammo, Cymbush, Cynoff, Cyperkill, Fenom, Arrivo	Pyrethroid Type II	Sodium Channel Agonist	ICI Americas (Zeneca) Goldsboro, NC
15	DDT	DDT	Chlorinated Hydrocarbon	Sodium Channel Agonist	Synthesis by B.D. Hammock, UC Davis, CA
20	Endosulfan	Thiodan, Cyclodan, Thiosulfan, Malix, Thionex, Tiovel	Cyclodiene	GABA Channel Agonist	FMC Corp. Ag Chemical Group Middleport, NY
25	Methomyl	Lannate, Nudrin	Carbamate	ACHe Inhibitor	E.I. DuPont de Nemours & Co. Wilmington, DE
30	Profenofos	Curacon, Polycron, Selecron	Organo- phosphate	ACHe Inhibitor	Ciba-Geigy Greensboro, NC

---

GABA, g-aminobutyric acid; AChE, acetylcholinesterase

TABLE 3B

Time-Response of Neonate Larvae of Tobacco Budworm to Recombinant or Wild-Type AcNPV with Low Concentrations of Classical Insecticides

	Treatment <sup>a</sup>	n	Slope $\pm$ SE	LT <sub>10</sub> (h; 95% CI)	LT <sub>50</sub> (h; 95% CI)	LT <sub>90</sub> (h; 95% CI)
5	Wt AcNPV	204	9.26 $\pm$ 0.41	64.9 (59.0 - 69.7)	89.8 (85.0 - 93.4)	123 (115 - 134)
	AcAaIT	276	7.73 $\pm$ 0.25	47.9 (44.8 - 50.6)	70.2 (68.0 - 72.4)	103 (98.3 - 109)
	AcJHE.KK	90	7.14 $\pm$ 0.53	50.5 (24.1 - 62.7)	76.3 (60.8 - 99.7)	115 (91.5 - 266)
	Allethrin	114	1.46 $\pm$ 0.19	17.3 (11.5 - 22.5)	132 (106 - 185)	1000 (552 - 2680)
10	Allethrin + AcAaIT	125	3.54 $\pm$ 0.20	17.5 (12.6 - 21.7)	40.2 (34.9 - 45.5)	92.5 (78.2 - 118)
	Allethrin + WtAcNPV	119	2.31 $\pm$ 0.19	20.0 (16.1 - 23.5)	71.9 (65.5 - 80.0)	258 (205 - 354)
	Cypermethrin	110	1.49 $\pm$ 0.24	17.1 (7.99 - 24.8)	124 (97.5 - 195)	902 (436 - 4190)
15	Cypermethrin + AcAaIT	112	3.19 $\pm$ 0.27	12.5 (5.60 - 18.5)	31.5 (22.7 - 38.1)	79.0 (65.3 - 111)
	Cypermethrin + WtAcNPV	109	2.09 $\pm$ 0.23	16.0 (10.9 - 20.7)	65.8 (59.3 - 73.8)	270 (203 - 418)
	Cypermethrin + AcJHE.KK	83	3.81 $\pm$ 0.27	24.2 (18.7 - 28.8)	52.5 (47.3 - 58.0)	114 (97.8 - 142)
20	DDT	80	1.45 $\pm$ 0.30	12.6 (3.90 - 23.8)	96.7 (73.5 - 165)	740 (318 - 10320)
	DDT + AcAaIT	80	3.12 $\pm$ 0.32	16.0 (7.84 - 23.0)	41.2 (31.4 - 48.7)	107 (84.6 - 160)
	DDT + WtAcNPV	80	2.25 $\pm$ 0.29	17.7 (5.78 - 28.2)	66.0 (49.9 - 84.6)	245 (159 - 688)
	Endosulfan	80	1.51 $\pm$ 0.27	10.4 (3.75 - 17.3)	73.3 (61.0 - 90.3)	514 (296 - 1580)
25	Endosulfan + AcAaIT	80	4.23 $\pm$ 0.35	23.8 (16.5 - 29.7)	47.7 (40.6 - 54.2)	95.4 (81.6 - 121)
	Endosulfan + WtAcNPV	80	2.57 $\pm$ 0.30	24.7 (14.5 - 33.0)	78.1 (66.7 - 93.6)	247 (176 - 460)
	Methomyl	125	0.69 $\pm$ 0.20	4.84 (0.12 - 13.1)	358 (182 - 4070)	26563 (2870-1.25x10 <sup>6</sup> )
30	Methomyl + AcAaIT	125	3.27 $\pm$ 0.28	13.7 (9.16 - 17.8)	34.3 (29.2 - 38.8)	85.7 (73.3 - 108)
	Methomyl + WtAcNPV	125	2.37 $\pm$ 0.22	14.6 (8.34 - 20.2)	50.3 (42.5 - 58.1)	173 (132 - 276)
	Profenofos	150	0.76 $\pm$ 0.19	8.10 (2.61 - 17.5)	401 (260 - 676)	19810 (9472-1.88x10 <sup>4</sup> )
35	Profenofos + AcAaIT	150	3.81 $\pm$ 0.23	24.5 (20.3 - 28.2)	53.2 (49.2 - 57.1)	115 (103 - 133)
	Profenofos + WtAcNPV	150	2.33 $\pm$ 0.19	16.5 (10.8 - 21.6)	58.5 (52.0 - 65.2)	208 (163 - 302)

<sup>a</sup> Dose-response curves were developed for the classical insecticides using neonate larvae of *H. virescens*. Neonate larvae infected with NPVs were exposed to 2000 polyhedrin inclusion bodies (PIBs;  $\geq 10^6$ ) using a droplet feeding assay. Three NPVs were utilized including WtAcNPV, AcAaIT, and AcJHE.KK. Larvae which were to be combined with an insecticide were immediately transferred to vials treated with a LC<sub>10</sub>-LC<sub>20</sub> (24 h) of the insecticide. Mortality was recorded at 8 to 12 h intervals, and the data were analyzed using a log-probit analyses program (POLO-PC).

40

However, as shown by the data of Table 5, the combination of the recombinant AcAaIT with low rates of either pyrethroid produced a dose-response greater than an additive response. Thus, we observed that AcAaIT in combination with allethrin or cypermethrin resulted in 30.0 and 38.7% reductions in the  $LT_{50}$ s, respectively, when compared to AcAaIT alone. The  $LT_{50}$ s for the combination of AcAaIT and allethrin or cypermethrin were 40.2 h and 31.5 h post infection, respectively (Table 3). Both of these  $LT_{50}$ s are lower than the theoretical (additive effect)  $LT_{50}$ s of 45.8 h and 44.8 h post infection, respectively. These data suggest that the pyrethroids and AaIT interacted in a manner, which magnifies the rate of kill beyond potentiation and are synergizing (supplemental synergism - quicker effect than the algebraic sum of the single effects) the rate of kill by AcAaIT.

Another recombinant virus, AcJHE.KK, was used as a control for the recombinant NPV, AcAaIT. AcJHE.KK expresses a modified version of juvenile hormone esterase, an insect-derived enzyme important in the regulatory development of many lepidopterous insects. The modified JHE has been shown to be insecticidal to several lepidopterous insects (Bonning and Hammock, 1994). Although the mode of action of JHE.KK is not yet fully understood, we can assume it does not interfere with the nervous system of insects. We found a 23.8% reduction in the  $LT_{50}$  when AcJHE.KK was combined with cypermethrin. The  $LT_{50}$  for this combination was 52.5 h which is significantly higher than 47.2 h predicted for the theoretical combination and indicates a subadditive not a supplemental effect (Table 5). Since AcJHE.KK has a  $LT_{50}$  similar to that of AcAaIT these data indicate that the pyrethroids do not synergize indiscriminately any baculovirus with enhanced speed of kill. Thus, these data provide further support for a supplemental



synergistic relationship between the recombinant virus, ACAAiT, and pyrethroid insecticides.

As further shown by Table 3B, another sodium channel agonist, DDT, was also screened for interactions with the NPVs. A combination of DDT and wild-type AcNPV resulted in a 24% reduction in the  $LT_{50}$  (66.0 h) when compared to wild-type AcNPV alone (Table 3B). Our data indicate that this combination produced a subadditive effect (46.7 h, theoretical). DDT combined with ACAAiT resulted in a 29.0% reduction in the  $LT_{50}$  (41.2 h) when compared to ACAAiT alone. The results of this combination suggest a purely additive response (theoretical  $\approx$  actual) since the theoretical time has a very similar  $LT_{50}$  of 40.8 hours.

Yet another insecticide, endosulfan (a cyclo-diene), was also combined with the NPVs to explore any interactions. Interestingly, our data indicate an antagonistic interaction when endosulfan was combined with the wild-type AcNPV. The  $LT_{50}$  for the combination was 78.1 h which was later than the 73.3 h produced by the insecticide alone indicating an antagonistic response. However, the combination of ACAAiT with endosulfan resulted in an  $LT_{50}$  of 47.7 h, which is significantly quicker than either ACAAiT or endosulfan alone and indicates a subadditive response.

Yet additional studies were performed with a carbamate (methomyl) or an organophosphate (profenofos) insecticide in combination with AcNPV or ACAAiT. Interestingly, both insecticide and wild-type AcNPV combinations resulted in supplemental synergism. The derived  $LT_{50}$  for methomyl and wild-type AcNPV was 50.3 h compared to 71.9 h for the theoretical response (Table 3). Likewise, profenofos in combination with wild-type AcNPV resulted in a  $LT_{50}$  of 58.5 h versus 73.5 h for the theoretical additive response. Similarly, the combinations of methomyl or profenofos with ACAAiT resulted in

supplemental synergistic responses. The  $LT_{50}$ s for methomyl or profenofos with AcAaIT were 34.3 h and 53.2 h, respectively. The combination of AcAaIT and methomyl show a 61.8% quicker rate of kill than wild-type AcNPV alone, and a 36.0% quicker rate than AcAaIT alone. Profenofos and AcAaIT result in a 40.8 and 17.0% reduction in time of kill compared to wild-type AcNPV and AcAaIT, respectively.

Overall, our data indicate that the pyrethroid insecticides and ACHE inhibitors appear to provide the greatest chance of synergizing the recombinant virus, AcAaIT, in the field and for this reason, use of the recombinant AcAaIT is particularly preferred. However, other recombinant strains of baculovirus expressing the AaIT gene would be also preferred (i.e. HZAaIT or SfAaIT). For the pyrethroids these data suggest that a simultaneous exposure of AaIT and pyrethroid may result in a synergistic interaction at the insect sodium channel. Based on these results, varying rates of pyrethroids in combination with AcAaIT or other recombinant viruses expressing AaIT should be tested for efficacy in the field. The low rates of pyrethroids we are suggesting should have minimal effect on beneficial arthropods, thus not compromising the host specificity provided by the virus. The combination is believed also useful in resistance management strategies.

Both carbamate and organophosphate insecticides act as poisons at the peripheral nervous system. These compounds inhibit the enzyme, acetylcholinesterase, resulting in an accumulation of acetylcholine in the synapse regions of nerves. This accumulation of acetylcholine produces a plethora of neurological responses resulting in a paralysis of the insect. Therefore, it could be suggested that the simultaneous

action of the insecticide and the peptide toxin would result in a supplemental synergistic response. It is difficult to account for the contribution, if any, made by the virus itself. However, NPVs can infect nerve cells, and eventually the infection results in lysis of the cell. One could envision that the lysis of nerve cells by NPV infection would result in an increase of neurotransmitters in the peripheral nervous system which could lead to neuromuscular abnormalities. This action might account for the supplemental synergism observed with both wild-type and recombinant NPVs when combined with methomyl or profenofos.

As indicated previously, these studies have emphasized the effects of combinations of viral and chemical insecticides on  $LT_{50}$ , but the same approaches could be applied to dose as well as time. We used here approximate  $LD_{10S}$  (24 h) of the pesticides reported here to minimize the symptomology and mortality resulting directly from the chemical pesticide (Table 4). In practice a variety of factors would influence the rates used, including the cost and availability of the biological and chemical pesticide, environmental concerns, level of resistant populations in the field, and other pest management considerations.

25

#### EXAMPLE 5

In this study we compared the LTs of wild-type and AcAaIT against pyrethroid-susceptible and -resistant (PEG) larvae of *H. virescens*. The response (LTs) of the susceptible and PEG neonate larvae to wild-type AcNPV and AcAaIT were very comparable. Statistical analysis (POLO-PC) of the data indicated no significant difference in the slopes (PEG = 9.51, susceptible = 9.82, 95% CL) or intercepts. However, when these viruses were tested for effectiveness against

pyrethroid-resistant larvae our results lead to the surprising results as follows. Analysis of the data indicated a significant difference between the slopes (PEG = 13.9, susceptible = 7.73, 95% CL) and intercepts for wild-type and AcAaIT, but our results indicated that the resistant strain, not the susceptible strain, was killed in a more timely manner. The  $LT_{90}$  for the Stoneville strain was 103 h post infection versus 78.1 h for the PEG strain representing a highly significant 24.1% quicker rate of kill against the resistant strain.

These results indicate that a recombinant baculovirus such as AcAaIT may be more effective against pyrethroid-resistant larvae than pyrethroid-susceptible larvae of *H. virescens* in field situations and could be used as an effective agent to combat pyrethroid resistance.

Thus, one aspect of the invention is use of AcAaIT to control outbreaks of pest insects resistant to pyrethroids. Furthermore, AcAaIT could be integrated into a resistance management strategy to delay or prevent pyrethroid resistance in the field. In addition to reducing resistant populations, this strategy should extend the efficacy of the pyrethroid insecticides.

25

#### EXAMPLE 6

We investigated the interaction of low rates ( $LC_{10}$ - $LC_{20}$  at 24 h, Table 4) of several insecticides in combination with wild-type AcNPV or AcAaIT ( $>LC_{99}$ ) in neonate larvae of *H. virescens*. Two of the compounds studied were pyrethroids, allethrin (Type I pyrethroid) or cypermethrin (Type II pyrethroid). In our study we found a subadditive response when low rates of allethrin or cypermethrin were combined with wild-type AcNPV.

TABLE 4

	Compound	n	$\mu\text{g}$ Insecti- cide/Vial	% Mortality at 24 hours
	Allethrin	121	0.05	14.0
5	Cypermethrin	110	0.008	11.8
	DDT	80	0.02	18.8
	Endosulfan	80	0.002	17.5
	Methomyl	125	0.02	20.8
10	Profenofos	150	0.06	16.3

Lethal concentrations of insecticides ( $\text{LC}_{10}$ - $\text{LC}_{20}$ ) were chosen based on results from Campanhola & Plapp (1989).

Referring to Table 5 below, a combination of allethrin and wild-type AcNPV reduced the  $\text{LT}_{50}$  to 71.9 h (53.1 h, theoretical additive effect) which is a 17.9% reduction when compared to wild-type AcNPV alone. Likewise, cypermethrin in combination with wild-type AcNPV produced a  $\text{LT}_{50}$  of 65.8 h (52.1 h, theoretical) which represents a 24.2% reduction when compared to wild-type AcNPV. These results suggest that the pyrethroids and the wild-type AcNPV act independently of one another but result in a subadditive response since the combination is positive although it is significantly less than a theoretical interaction (additive effect).

The combination of the recombinant NPV, AcAaIT, with low rates of either pyrethroid produced a dose-response greater than an additive response. We observed that AcAaIT in combination with allethrin or cypermethrin resulted in 30.0 and 38.7% reductions in the  $\text{LT}_{50}$ s, respectively, when compared to AcAaIT alone. The  $\text{LT}_{50}$ s for the combination of AcAaIT and allethrin or cypermethrin were 40.2 h and 31.5 h. post infection, respectively (Table 5). Both of these  $\text{LT}_{50}$ s are lower than the theoretical (additive effect)  $\text{LT}_{50}$ s of 45.8 h and 47.2 h post infection, respectively. These data illustrate the synergistic effect.

TABLE 5

Interactions of Recombinant or  
Wild-Type AcNPV with Low Concentrations of Classical Insecticides

	Treatment <sup>a</sup>	Actual Kill Time (LT <sub>50</sub> , h)	Theoretical Kill Time (h) <sup>b</sup>	% Reduction from WtAcNPV	% Reduction from RecNPV	Effect <sup>c</sup>
5	Wt AcNPV	70.2	---	21.8	---	---
	AcAaIT	89.8	---	---	---	---
	AcJHE.KK	76.2	---	15.0	---	---
	Allethrin	132	---	---	---	---
10	Allethrin + AcAaIT	40.2	45.8	55.2	30.0	Supplemental
	Allethrin + WtAcNPV	71.9	53.8	17.9	---	Subadditive
	Cypermethrin	124	---	---	---	---
15	Cypermethrin + AcAaIT	31.5	44.8	64.9	38.7	Supplemental
	Cypermethrin + AcJHE.KK	65.8	52.1	24.2	---	Subadditive
	Cypermethrin + WtAcNPV	52.5	47.2	41.2	23.8	Subadditive
20	DDT	96.7	---	---	---	---
	DDT + AcAaIT	41.2	40.8	54.1	29.0	Additive
	DDT + WtAcNPV	66.0	46.7	24.0	---	Subadditive
	Endosulfan	73.3	---	---	---	---
25	Endosulfan + AcAaIT	47.7	36.0	46.9	22.6	Subadditive
	Endosulfan + WtAcNPV	78.1	40.7	11.9	---	Antagonism
	Methomyl	358	---	---	---	---
30	Methomyl + AcAaIT	34.3	59.5	61.8	36.0	Supplemental
	Methomyl + WtAcNPV	50.3	71.9	39.5	---	Supplemental
35	Profenofos	401	---	---	---	---
	Profenofos + AcAaIT	53.2	59.9	40.8	17.0	Supplemental
	Profenofos + WtAcNPV	58.5	73.5	31.3	---	Supplemental

40 LT = lethal time; RecNPV = Recombinant NPV.

- ° Dose-response curves were developed for the classical insecticides using neonate larvae of *H. virescens*. Larvae infected with NPVs were exposed to 2000 polyhedrin inclusion bodies (PIBs;  $>LC_{90}$ ) using a droplet feeding assay. Larvae were immediately transferred to vials treated with an approximate  $LC_{10}$ - $LC_{20}$  (24 h) of the insecticide and scored every 8 to 12 h.
  - 5    ° Theoretical Kill Time =  $1 + [\text{Reciprocal of } LT_{50} \text{ for virus alone} + \text{Reciprocal of } LT_{50} \text{ for insecticide alone}]$ .
  - ° Effect - Supplemental synergism is Actual Time  $>$  Theoretical Time; Additive synergism is Actual Time = Theoretical Time; Subadditive Synergism is Actual Time  $<$  Theoretical Time and Actual Time  $>$  Fastest Time Alone; Antagonism is Actual Time  $<$  Fastest Time Alone.
- 

10                    The foregoing examples illustrate certain embodiments of the present invention, and are not intended to limit the scope of the invention, which is defined in the appended claims.

It is Claimed:

1. A method for controlling pests from the group insects, acarids, and nematodes comprising:  
treating said pests or their loci with a combination of recombinant insect virus and organic  
5 insecticide.
2. The method as in claim 1 wherein the recombinant insect virus expresses a foreign protein or a functional derivative thereof in pest cells infected therewith.
3. The method as in claim 2 wherein the expressed foreign protein or functional derivative is an insect toxin.
4. The method as in claim 1 wherein the insect virus is a baculovirus.
5. The method as in claim 4 wherein the baculovirus is a nuclear polyhedrosis virus.
6. The method as in claim 5 wherein the baculovirus is from *Autographa californica*, *Anagrapha falcifera*, *Anticarsia gemmatilis*, *Buzura suppressoria*, *Cydia pomonella*, *Heliocoverpa zea*, *Heliothis arrigera*,  
5 *Mariestia brassicae*, *Plutella xylostella*, *Spodoptera exigua*, *Spodoptera littoralis*, or *Spodoptera litura*.
7. The method as in claim 1 or 5 wherein the organic insecticide is a sodium channel agonist.
8. The method as in claim 7 wherein the sodium channel agonist is a pyrethroid.



9. The method as in claim 1 or 5 wherein the organic insecticide is an acetylcholinesterase inhibitor.
10. The method as in claim 9 wherein the acetylcholinesterase inhibitor is an organophosphate.
11. The method as in claim 9 wherein the acetylcholinesterase inhibitor is a carbamate.
12. The method as in claim 1 or 5 wherein the organic insecticide is a sodium channel blocking agent.
13. The method as in claim 12 wherein the sodium channel blocking agent is pyrazoline.
14. The method as in claim 1 or 5 wherein the organic insecticide is a nicotinic acetylcholine binding agent.
15. The method as in claim 1 or 5 wherein the organic insecticide is a gabaergic binding agent.
16. The method as in claim 1 or 5 wherein the organic insecticide is an octapamine agonist or antagonist.
17. The method as in claim 1 or 5 wherein the organic insecticide is an oxphos uncoupler.
18. The method as in claim 1 or 5 wherein the organic insecticide is an agent that binds to a cation channel.
19. The method as in claim 18 wherein the organic insecticide is a peptide.

20. A method of controlling insecticide resistant pests comprising:

treating said pests or their loci with a recombinant insect virus in conjunction with applying to  
5 the loci thereof an amount of insecticide to which the pest is resistant.

21. The method as in claim 20 wherein the amount of insecticide is reduced with respect to a determinable killing dose of insecticide when applied in the absence of the virus.

22. The method as in claim 20 wherein the insect virus is a baculovirus.

23. A method of treating insecticide resistant pests comprising:

determining a first amount of insecticide to which the pest is resistant, said first amount being  
5 sufficient to kill at least about half of the insects; and

administering a predetermined amount of said insecticide to the loci of said pests in conjunction with a recombinant baculovirus, the second amount being  
10 lower than said first dose of the determining step.

24. The method as in claim 20 or 23 wherein the insecticide is a pyrethroid.

25. The method as in claim 20, 23, or 24 wherein the baculovirus is a nuclear polyhedrosis virus.

26. The method as in claim 24 wherein the nuclear polyhedrosis virus is *Autographa californica*.

27. The method as in claim 25 wherein the recombinant baculovirus expresses an insect-selective toxin.

28. The method as in claim 26 wherein the toxin is from *Androctonus australis*.

29. The method as in claim 27 wherein the pest is of the genera *Spodoptera*, *Trichoplusia*, or *Heliothis*.

30. The method as in claim 27 wherein the pest is *Heliothis virescens* or *Heliocoverpa zea*.

31. The method as in claim 29 wherein the insecticide is a pyrethroid.

- 34 -

## AMENDED CLAIMS

[received by the International Bureau on 30 October 1995 (30.10.95);  
original claims 1-3,17,20,21,23,25,27 amended;  
new claims 32-35 added; remaining claims unchanged (5 pages)]

1. A method for controlling pests from the group insects, acarids, and nematodes comprising:  
treating said pests or their loci with a  
5 combination of recombinant insect virus expressing a foreign protein in cells of said pests infected therewith, and an organic insecticide, the combination effective to increase speed of pest kill.
2. The method as in claim 1 wherein the  
10 foreign protein is an insect toxin or a functional derivative of the insect toxin that is active against said pests.
3. The method as in claim 2 wherein the insect toxin is a scorpion toxin.
- 15 4. The method as in claim 1 wherein the insect virus is a baculovirus.
5. The method as in claim 4 wherein the baculovirus is a nuclear polyhedrosis virus.
- 20 6. The method as in claim 5 wherein the baculovirus is from *Autographa californica*, *Anagrapha falcifera*, *Anticarsia gemmatalis*, *Buzura suppressoria*, *Cydia pomonella*, *Heliocoverpa zea*, *Heliothis arrigera*, *Mariestia brassicae*, *Plutella xylostella*, *Spodoptera exigua*, *Spodoptera littoralis*, or *Spodoptera litura*.
- 25 7. The method as in claim 1 or 5 wherein the organic insecticide is a sodium channel agonist.

AMENDED SHEET (ARTICLE 19)

- 35 -

8. The method as in claim 7 wherein the sodium channel agonist is a pyrethroid.

9. The method as in claim 1 or 5 wherein the organic insecticide is an acetylcholinesterase inhibitor.  
5

10. The method as in claim 9 wherein the acetylcholinesterase inhibitor is an organophosphate.

11. The method as in claim 9 wherein the acetylcholinesterase inhibitor is a carbamate.

10 12. The method as in claim 1 or 5 wherein the organic insecticide is a sodium channel blocking agent.

13. The method as in claim 12 wherein the sodium channel blocking agent is pyrazoline.

15 14. The method as in claim 1 or 5 wherein the organic insecticide is a nicotinic acetylcholine binding agent.

15. The method as in claim 1 or 5 wherein the organic insecticide is a gabaergic binding agent.

20 16. The method as in claim 1 or 5 wherein the organic insecticide is an octapamine agonist or antagonist.

17. The method as in claim 1 or 5 wherein the organic insecticide is an oxidative phosphorylation uncoupler.

-36 -

18. The method as in claim 1 or 5 wherein the organic insecticide is an agent that binds to a cation channel.

19. The method as in claim 18 wherein the organic insecticide is a peptide.

20. A method of controlling insecticide resistant pests comprising:

treating said pests or their loci with a recombinant insect virus expressing a foreign protein in cells of said pests infected therewith, in conjunction with applying to the loci thereof an amount of insecticide to which said pest is resistant, the amount of insecticide applied being not greater than the dose that would have been applied for nonresistant pests, the recombinant insect virus and the insecticide effective together to increase speed of kill for said pest.

21. The method as in claim 20 wherein the speed of kill is faster for said pests than for the nonresistant pests.

22. The method as in claim 20 wherein the insect virus is a baculovirus.

23. A method of treating insecticide resistant pests comprising:

determining a first amount of insecticide to which the pest is resistant, said first amount being sufficient to kill at least about half of the insects; and

administering a second amount of said insecticide to the loci of said pests in conjunction with a recombinant baculovirus expressing a foreign protein in cells of said pests infected therewith, the

- 37 -

second amount being lower than said first amount of the determining step and the speed of kill being faster for the insecticide and the recombinant baculovirus than for the insecticide or the baculovirus alone.

5                   24. The method as in claim 20 or 23 wherein the insecticide is a pyrethroid.

                  25. The method as in claim 23 wherein the baculovirus is a nuclear polyhedrosis virus.

10                   26. The method as in claim 24 wherein the nuclear polyhedrosis virus is *Autographa californica*.

                  27. The method as in claim 25 wherein the foreign protein is an insect-selective toxin.

                  28. The method as in claim 26 wherein the toxin is from *Androctonus australis*.

15                   29. The method as in claim 27 wherein the pest is of the genera *Spodoptera*, *Trichoplusia*, or *Heliothis*.

                  30. The method as in claim 27 wherein the pest is *Heliothis virescens* or *Heliocoverpa zea*.

20                   31. The method as in claim 29 wherein the insecticide is a pyrethroid.

                  32. The method as in claim 1 or 5 wherein speed of pest kill is faster than that set by a theoretical combination kill time, C, wherein C is defined by the relationship:

$$1/C = 1/A + 1/B$$

AMENDED SHEET (ARTICLE 19)


- 38 -

wherein A is the individual kill time at a particular concentration of recombinant insect virus and B is the individual kill time at a particular concentration of organic insecticide.

- 5                   33. The method as in claim 5 wherein the baculovirus is the nuclear polyhedrosis virus *Autographa californica* expressing the toxin from *Androctonus australis* and the insecticide is a pyrethroid or an acetylcholinesterase inhibitor.
- 10                   34. The method as in claim 20 wherein the insect virus is the nuclear polyhedrosis virus *Autographa californica* expressing a toxin from *Androctonus australis* and the insecticide is a pyrethroid.
- 15                   35. The method as in claim 23 wherein the baculovirus is the nuclear polyhedrosis virus *Autographa californica* expressing a toxin from *Androctonus australis* and the insecticide is a pyrethroid.



FIG. 1



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/06664

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 63/00; 27/00; C12N 7/01

US CL : 424/93.2, 93.6, 199.1, 204.1, 405; 435/235.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.2, 93.6, 199.1, 204.1, 405

435/172.3, 235.1, 240.2, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,668,511 (ASPIROT <i>ET AL.</i> ) issued 26 May 1987, see entire document.	1-31
Y	US, A, 5,238,724 (BJOSTAD <i>ET AL.</i> ) issued 24 August 1993, see entire document.	1-31
Y	US, A, 5,162,308 (BROWN <i>ET AL.</i> ) issued 10 November 1992, see entire document.	1-31
Y	US, A, 4,888,340 (NEH <i>ET AL.</i> ) issued 19 December 1989, see entire document.	1-31
Y	BIO/TECHNOLOGY, Volume 9, issued September 1991, McCutchen et al., "Development Of A Recombinant Baculovirus Expressing An Insect-Selective Neurotoxin: Potential For Pest Control", pages 848-852, see entire document.	1-31



Further documents are listed in the continuation of Box C.



See patent family annex.

\*

Special categories of cited documents:

\*A\*

document defining the general state of the art which is not considered to be of particular relevance

\*E\*

earlier document published on or after the international filing date

\*L\*

document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\*

document referring to an oral disclosure, use, exhibition or other means

\*P\*

document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\*

document member of the same patent family

Date of the actual completion of the international search

03 JULY 1995

Date of mailing of the international search report

28 AUG 1995

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHRISTOPHER S. F. LOW

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/06664

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Science, Volume 261, issued 30 July 1993, Moffat, "New Chemicals Seek To Outwit Insect Pests", pages 550-551, see entire document.	1-31

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/06664

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Automated Patent System - USPAT and JPOABS

DIALOG - BIOSIS

Search Terms: insecticide, pesticide, baculovirus, toxin, pyrethrin, carbamate, organophosphorus, pyrazoline, imidacloprid, formamidine, Autographica,